



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2019

Furin inhibition prevents hypoxic and TGF -mediated blood-brain barrier disruption

Baumann, Julia ; Huang, Sheng-Fu ; Gassmann, Max ; Tsao, Chih-Chieh ; Ogunshola, Omolara O

Abstract: Hypoxic blood-brain barrier (BBB) dysfunction is a common feature of CNS diseases however mechanisms underlying barrier disturbance are still largely unknown. This study investigated the role of transforming growth factor (TGF), a cytokine known to induce expression of the proprotein convertase Furin, in hypoxia-mediated barrier compromise. We show that exposure of brain endothelial cells (ECs) to hypoxia (1% O₂) rapidly stimulates their migration. Additional exogenous TGF (0.4nM) exposure potentiated this effect and increased Furin expression in a TGF type I receptor activin-like kinase 5 (ALK5) - dependent manner (prevented by 10 M SB431542). Furin inhibition prevented hypoxia-induced EC migration and blocked TGF -induced potentiation suggesting existence of a feedback loop. TGF and Furin were also critical for hypoxia-induced BBB dysfunction. TGF treatment aggravated hypoxia-induced BBB permeability but ALK5 or Furin blockade reversed injury-induced permeability changes. Thus during insult Furin compromises endothelial integrity by mediating the effects of TGF. Targeting the Furin or ALK5 pathway may offer novel therapeutic strategies for improving BBB stability and CNS function during disease.

DOI: <https://doi.org/10.1016/j.yexcr.2019.111503>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-172353>

Journal Article

Accepted Version

Originally published at:

Baumann, Julia; Huang, Sheng-Fu; Gassmann, Max; Tsao, Chih-Chieh; Ogunshola, Omolara O (2019). Furin inhibition prevents hypoxic and TGF -mediated blood-brain barrier disruption. *Experimental Cell Research*, 383(2):111503.

DOI: <https://doi.org/10.1016/j.yexcr.2019.111503>

Accepted Manuscript

Furin inhibition prevents hypoxic and TGF β -mediated blood-brain barrier disruption

Julia Baumann, Sheng-Fu Huang, Max Gassmann, Chih-Chieh Tsao, Omolara O. Ogunshola



PII: S0014-4827(19)30344-1

DOI: <https://doi.org/10.1016/j.yexcr.2019.111503>

Article Number: 111503

Reference: YEXCR 111503

To appear in: *Experimental Cell Research*

Received Date: 5 April 2019


Revised Date: 25 June 2019

Accepted Date: 15 July 2019

Please cite this article as: J. Baumann, S.-F. Huang, M. Gassmann, C.-C. Tsao, O.O. Ogunshola, Furin inhibition prevents hypoxic and TGF β -mediated blood-brain barrier disruption, *Experimental Cell Research* (2019), doi: <https://doi.org/10.1016/j.yexcr.2019.111503>.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Proposed mechanism of Furin and TGF β at the BBB



Furin inhibition prevents hypoxic and TGF β -mediated blood-brain barrier disruption.

Running title: Furin-mediated BBB permeability

Julia Baumann, Sheng-Fu Huang, Max Gassmann, Chih-Chieh Tsao, and Omolara O. Ogunshola

Institute of Veterinary Physiology, Vetsuisse Faculty and Zurich Center Integrative Physiology (ZIHP), University of Zurich, Zurich, Switzerland.

Address correspondence to

Omolara O. Ogunshola, PhD

Institute of Veterinary Physiology, Vetsuisse Faculty, University of Zurich,
Winterthurerstrasse 260, CH-8057 Zurich, Switzerland

Tel: +41 44 635 8805, Fax: +41 44 635 8932, Email: larao@access.uzh.ch

ABSTRACT

Hypoxic blood-brain barrier (BBB) dysfunction is a common feature of CNS diseases however mechanisms underlying barrier disturbance are still largely unknown. This study investigated the role of transforming growth factor β (TGF β), a cytokine known to induce expression of the proprotein convertase Furin, in hypoxia-mediated barrier compromise. We show that exposure of brain endothelial cells (ECs) to hypoxia (1% O₂) rapidly stimulates their migration. Additional exogenous TGF β (0.4nM) exposure potentiated this effect and increased Furin expression in a TGF β type I receptor activin-like kinase 5 (ALK5) - dependent manner (prevented by 10 μ M SB431542). Furin inhibition prevented hypoxia-induced EC migration and blocked TGF β -induced potentiation suggesting existence of a feedback loop. TGF β and Furin were also critical for hypoxia-induced BBB dysfunction. TGF β treatment aggravated hypoxia-induced BBB permeability but ALK5 or Furin blockade reversed injury-induced permeability changes. Thus during insult Furin compromises endothelial integrity by mediating the effects of TGF β . Targeting the Furin or ALK5 pathway may offer novel therapeutic strategies for improving BBB stability and CNS function during disease.

Key words: Furin, ALK-5, barrier permeability, cell migration, primary endothelial cells

1 Introduction

2
3 A stable well-functioning blood-brain barrier (BBB) is crucial to maintain and control
4 cerebral homeostasis as well as ensure proper neuronal function. The BBB is formed
5 by specialized brain endothelial cells (EC) that limit paracellular movement of
6 substances to and from the brain parenchyma by their tight junctions (TJ) (1, 2).
7 Pericytes and perivascular astrocytes present at the vascular wall also contribute to
8 barrier maintenance (3). BBB dysfunction allows the influx of damaging molecules,
9 leading to neuronal cell loss and consequently cognitive decline and CNS disease (4).

10 Hypoxia is a key stress factor that induces BBB disturbance and increases vessel
11 permeability through a variety of different mechanisms including EC cytoskeletal
12 reorganization, TJ protein delocalization and altered molecular pathway signaling (5,
13 6). Hypoxic insults induce the expression of transforming growth factor- β 1 (7) and
14 during CNS injury and disease levels of bioactive TGF- β 1 increase (7-9).
15 Transforming growth factor betas (TGF- β s) are known multifunctional growth factors
16 that participate in regulation of key events during development, disease, and tissue
17 repair. In the brain TGF- β 1, 2 and 3 are broadly expressed with TGF- β 1 widely
18 recognized as an injury-related cytokine. Canonical TGF- β signaling is initiated by
19 ligand binding to a high-affinity transmembrane TGF- β type II receptor (TGF β RII)
20 that recruits and phosphorylates the type I receptors activin-like kinase 1 and 5
21 (ALK1, ALK5). Subsequent phosphorylation of the downstream effector proteins
22 Smad2 and Smad3 via ALK5 or Smad 1, 5 and 8 via ALK1 causes formation of a
23 heteromeric complex with Smad4, nuclear translocation and transcriptional
24 regulation (10). At the level of the endothelium TGF- β can signal through both
25 ALK1 and ALK5 and induce different molecular pathways depending on context or
26 dosage (11).

27 The pleiotropic cytokine TGF β -1 (henceforth referred to as TGF β) is involved in
28 many crucial physiological processes such as cell growth, cell differentiation,
29 migration and apoptosis (11) depending on the cell type. For example, whilst the
30 growth factor induced vessel formation in bovine capillary endothelial cells (BCEC),
31 others showed it promoted apoptosis in BCECs and HUVECs and could inhibit the
32 proliferation of bovine heart ECs (12). Targeted deletion of various TGF β signaling
33 pathway components in mice led to embryonic lethality due to impaired vascular
34 development with leaky vessels (13) and/or angiogenesis defects (14). In correlation
35 dysfunctional TGF β signaling has been observed in several pathological conditions
36 linked to vascular compromise such as cancer (15), fibrotic disease (16) and
37 atherosclerosis (17). The TGF β pathway is clearly a major regulator of blood vessel
38 characteristics.

39 TGF β has been shown to induce expression of the proprotein convertase Furin in
40 synovial cells and fibroblasts (18). Furin, a calcium-dependent ubiquitously expressed
41 cellular endoprotease (19), is known to process several protein precursors associated
42 with vascular remodeling and cell movement such as vascular endothelial growth
43 factor C (VEGF-C) (20), platelet derived growth factor (PDGF-BB) (21) and
44 membrane-type 1 matrix metalloprotease 1 (MT1-MMP) (22). Very recently it was
45 demonstrated that Furin deficiency in myeloid cells attenuated revascularization after
46 oxygen-induced retinopathy (23). Overall a crucial role for Furin in modulating
47 various vascular endothelia possibly via existence of feedback loops between the
48 enzyme and its substrates has been suggested. In this regard it is important to note that
49 a number of studies using cells of other origin have also convincingly shown Furin to

be a TGF β converting enzyme (24-26). Whether signaling between these two factors occurs at the BBB and/or modulates barrier dysfunction is unknown.

The aim of this study was to investigate the effect of TGF β on barrier stability during injury and the potential involvement of Furin in the process. We demonstrate that TGF β potentiates hypoxia-induced brain EC migration and BBB permeability, and that Furin plays a key role in mediating these effects.

Materials and Methods

Cell culture

The rat brain endothelial cell line RBE4 (27) was used for experiments at passages between 36 and 49 at 80-90% confluency. Cells were maintained on rat-tail collagen coated petri dishes in 1:1 α MEM/Ham's F-10 medium mixture supplemented with 10% FBS, 300 μ g/mL Geneticin purchased from Gibco[®] (Thermo Fisher Scientific, MA, USA) and 1ng/mL basic fibroblast growth factor (PeproTech, NJ, USA). Rat-tail collagen was isolated as previously described (28) and diluted to 250 μ g/mL in 70% Ethanol.

Primary endothelial cell isolation

Primary rat brain endothelial cells (primary ECs) were isolated from 8-10 week old male Wistar rats according to Coisne et al. (29), with slight modifications. Cortices were homogenized after removal of meninges in a Dounce homogenizer. Microvessels were isolated by adding an equal volume of 30% dextran solution to the homogenate and subsequent centrifugation at 3000g for 25 min at 4°C, then filtered through a 60 μ m nylon mesh to remove larger vessels. After digestion in HBSS buffer supplemented with 10mM HEPES, 0.1% BSA, 2mg/mL collagenase-dispase, 10 μ g/mL DNase I and 147ng/mL TLCK, for 45min at 37°C, microvessels were re-suspended in endothelial media (DMEM Glutamax II supplemented with 10% calf serum, BME amino acids, vitamin solution, 2mM L-glutamine, 1ng/mL bFGF, 59 μ g/mL gentamycin sulfate). After plating on collagen IV coated culture dishes pericyte contamination was prevented by exposing cultures to 3 μ g/mL puromycin-containing media for 24h. Cells were used in experiments on day 5 without passaging.

Hypoxic and substance exposures

Hypoxic exposures were performed in a hypoxic glove box chamber at 37°C, 5% CO₂ and 1% O₂ (InVivoO₂ 400, Ruskinn Technologies, Pencoe UK). An internal O₂ sensor monitored O₂ concentrations constantly. The following substances were added exogenously to the cell media: 0.4nM human TGF β -1 (PeproTech) dissolved in 10mM citric acid, 10 μ M ALK5 inhibitor SB431542 (Tocris, Bristol, UK) in DMSO, 20 μ M Naphthofluorescein (Furin inhibitor, Santa Cruz, TX, USA) in DMSO, 20 μ M Decanoyl-Arg-Val-Lys-Arg-chloromethylketone (CMK, Furin inhibitor, Santa Cruz) in H₂O. Cells were pretreated with inhibitors for 1h prior to hypoxic and TGF β exposures.

Cell viability

Viability of confluent cells was assessed using MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide). MTT solution (Sigma-Aldrich, MO, USA) was added to the media (final concentration 0.5mg/mL) and incubated for 1h at 37°C. The

media was removed and formazan crystals dissolved in DMSO. Optical density was measured at 570nm on a spectrophotometer (Thermo Labsystems, Multiskan RC Model 351) with reference filter at 670nm.

Cell migration

RBE4 cells were grown in 6-well plates and a scratch was introduced vertically with a 200 μ L pipette tip. Three pictures were taken per well before and after the exposures with an inverted microscope coupled to an 8-bit CCD camera (Axiocam HR, Carl Zeiss, Switzerland). Image analysis using ImageJ software (NIH, USA) was used to determine scratch width. Mean migration (μ m/h) was calculated by evaluating the difference in gap size before and after the exposure.

Permeability assay

ECs were grown to confluency on 0.4 μ m Transwells (Corning Incorporated, NY, USA). After exposures fresh medium containing 1mg/mL Lucifer yellow CH, lithium salt (Life technologies, OR, USA) was added to the upper compartment. At 9, 15, 30, 45 and 60 min aliquots were taken from the bottom compartment and tracer flux was determined with a fluorescence plate reader (FLx800, Biotek Instruments, Winooski, VT). A clearance slope established from the measurements obtained at the different time points was used to calculate the permeability coefficient (Pe) (30).

Western blotting

Cells were lysed in whole cell lysis buffer (50mM Tris, 150mM NaCl, 1% Triton X-100, 1% NP-40) supplemented with protease inhibitor cocktail (Calbiochem, Darmstadt, Germany), 1mM sodium orthovanadate and 0.5mM phenylmethanesulfonyl fluoride. Protein concentrations were determined with a Pierce BCA protein assay (Thermo Fischer Scientific). Proteins (50 μ g) were separated by SDS-PAGE then transferred to nitrocellulose membranes and blocked with 5% non-fat dried milk or 5% BSA in TBS. After incubation in primary antibodies over night at 4°C membranes were washed then incubated with the corresponding HRP-conjugated secondary antibody. Bands were visualized with a luminescent image analyzer (Fujifilm, LAS-3000, Switzerland) and quantification was performed with ImageJ software (NIH, USA) using β -actin as loading control. The following antibodies were used: anti-Furin (SC-20801, Santa Cruz), anti-HIF-1 α (NB100-479 W-1, NovusBio), anti-pSmad2 (138D4, Cell Signaling), anti-Smad2 (D43B4, Cell Signaling), anti-Smad1/5 (D5B10, Cell Signaling), anti-Smad1 (D59D7, Cell Signaling) and β -actin (A5441, Sigma-Aldrich).

Quantitative RT-PCR

RNA was isolated from cells using TRIzol[®] reagent (Thermo Fisher). Total RNA (1 μ g per sample) was reverse transcribed with oligo-dT primers using the ImProm-II Reverse Transcriptase kit (Promega, Switzerland) according to the manufacturers protocol. Real-time quantitative PCR (qPCR) was performed with an ABI 7500 Fast Real-Time PCR System using Power SYBR[®] Green Master Mix (Applied Biosystems, Switzerland). The following primers were employed: Furin 5'-CAG AAG CAT GGC TTC CAC AAC-3' and 5'-TGT CAC TGC TCT GTG CCA GAA-3' and β -actin 5'-CTG GCT CCT AGC ACC ATG AAG-3' and 5'-GCC ACC GAT CCA CAC AGA GT-3'. Serial cDNA dilution series were used to establish optimal primer conditions resulting in >90% efficiency. To exclude primer dimer and off-target amplifications melting curves were performed and confirmed by gel

electrophoresis. Data was normalized to β -actin and fold changes calculated using the $\Delta\Delta C_t$ method.

Statistical Analysis

Statistical Analyses were performed with Prism GraphPad 6. Results are shown as mean values \pm standard deviation with a minimum of 3 independent experiments. Statistical significance was determined by two-way ANOVA when comparing more than two different groups with different exposures or unpaired Student's t-test with homoscedasticity. A p value less than 0.05 was considered significant.

RESULTS

TGF β promotes EC migration

Till now the role of TGF β signaling has been mainly studied in peripheral ECs whereas its effects on the brain endothelium remain largely debated. In the first instance, we aimed to establish how increased TGF β levels impact brain EC quiescence during hypoxic injury using a classic scratch migration assay and the rat brain microvascular endothelial (RBE4) cell line. Notably, our studies have shown this cell line exhibits the same responses as primary cells but with increased sensitivity to oxygen deprivation (6). Scratch assays were performed during 6h normoxic and hypoxic exposure with exogenous TGF β and ALK5 inhibitor treatment (Fig. 1A&B). Quantification showed no significant changes under normoxic conditions with any of the treatments (Fig. 1C). In contrast hypoxia significantly increased EC migration and TGF β potentiated this effect. Notably, both hypoxic and TGF β -mediated migration were completely blocked when ECs were pretreated with the TGF β receptor I inhibitor (SB431542). Thus TGF β potentiates hypoxia-mediated cell migration via a TGF β R1 dependent mechanism.

Hypoxia and TGF β modulate Furin protein and expression and activation

Since existence of a feedback loop between TGF β and Furin has been implicated in various other (non-vascular) cells (24-26), we asked if a similar mechanism could underlie the movement we observed. As Furin mRNA levels were unaffected in our treatment conditions (Supp. Fig. 1) we evaluated whether Furin activation correlated with the TGF β -induced migratory effects. Hypoxic HIF-1 α stabilization confirmed the endogenous endothelial cell response was as expected and unaffected by TGF β treatment (Fig. 2A). Corresponding Furin immunoblots showed two prominent bands corresponding to the inactive precursor form at 96kDa and the mature active form at 90kDa (Fig. 2A). Quantitative analysis of the individual bands revealed a significant decrease of the Furin precursor under hypoxia. Notably, TGF β had no impact on normoxic precursor levels but induced hypoxic precursor levels by 50% to reach the baseline normoxic levels (Fig. 2B). Mature Furin levels remained unchanged by hypoxia compared to the normoxic baseline (Fig. 2C) but significantly increased during hypoxic TGF β exposure, with a trend to also increase in normoxia. This data suggests that during stress (hypoxic) conditions high TGF β levels might potentiate EC Furin activation by elevating precursor levels and inducing protein maturation.

Furin activation is dependent on ALK5 signaling

In ECs, canonical binding of TGF β to ALK5 causes downstream activation and phosphorylation of Smad2 resulting in stimulation of angiogenesis (31), whereas signaling via ALK1 induces phosphorylation of Smad1, 5 and 8 and subsequent endothelial quiescence (32). Exposure of brain ECs to TGF β but not hypoxia increased phosphorylated Smad2 levels whereas the opposite effect was observed on phospho-Smad1/5 levels (Supp. Fig. 2&3). In line with suppressing cell migration (Fig. 1), ALK5 blockade also prevented TGF β -mediated Smad2 phosphorylation but not the hypoxia-induced Smad1/5 activation. As Furin levels were also strongly potentiated by both hypoxia and TGF β (Fig. 2) we asked if ALK5 blockade would suppress these effects (Fig. 3). Indeed immunoblotting showed SB431542 abrogated expression of the Furin precursor and mature forms by more than 50% during hypoxia and TGF β exposure (Fig. 3). Interestingly, this pattern was also observed during normoxia (Supp. Fig. 4) suggesting Furin activity is highly dependent on ALK5 activation in both injury and physiological conditions.

Furin inhibition prevents EC migration

We next evaluated if Furin is the mediator of hypoxia- and TGF β -induced EC migration. A scratch assay was performed over 6h hypoxia as previously in the presence and absence of TGF β and the Furin inhibitors Naphthofluorescein (33) and CMK. Clear differences were apparent after visual inspection of the samples (Fig. 4A). Quantification showed hypoxia-induced cell migration was significantly potentiated when combined with exogenous TGF β as expected however treating cells with Furin inhibitors completely abrogated this response even in the presence of TGF β (Fig. 4B). Notably, both inhibitors also blocked normoxic EC migration by up to 60% (Supp. Fig. 5). These observations highlight that TGF β -induced EC migration is mediated by Furin and again underline an important physiological role for Furin.

TGF β potentiates hypoxia-mediated EC permeability

To understand whether injury-increased TGF β levels contribute to altered barrier integrity we performed permeability assays on primary microvascular ECs isolated from adult rat brain and RBE4 (data not shown) in presence and absence of exogenous TGF β (Fig. 5). Experiments were performed for 24h (Fig. 5A) and 48h (Fig. 5B) under normoxic and hypoxic conditions. Hypoxia significantly increased permeability of EC monolayers as frequently observed in many studies including our own (3). Exposure to TGF β further potentiated hypoxia-induced barrier dysfunction although no effects were observed under normoxia (Fig. 5A). Pretreatment with SB431542 completely abrogated TGF β and hypoxic driven effects to baseline normoxic levels. Indeed at both time points a similar outcome was seen although the effects at 48h were clearly more enhanced (Fig. 5A&B). Thus sustained injury-induced activation of the endothelial TGF β -ALK5 pathway culminates in barrier disturbance.

Furin inhibition prevents hypoxia-induced barrier permeability

Finally, due to the observed feedback between the proteins, we investigated if Furin inhibition would also suppress the barrier changes induced by hypoxic and/or TGF β exposure. Again we employed the transwell permeability assay using primary rat ECs (Fig. 6A) and RBE4 (data not shown). Blocking Furin activation with CMK was highly efficient at preventing hypoxia-mediated barrier disruption after 48h. Similar to observations in the migration assay, the CMK inhibitor resulted in a tighter monolayer than observed under normoxic conditions (Fig. 6A). Surprisingly however

the combination of CMK and exogenous TGF β dramatically increased EC barrier permeability by more than 5 fold. This led us to use MTT assays to assess if the observed permeability changes were due to compromised cell viability. Data showed that when combined with hypoxia TGF β treatment already compromised cell function up to 20%. Whilst CMK alone had no effect, the combination treatment reduced mitochondrial activity by 50% (Fig. 6B). Thus Furin blockade in the presence of elevated TGF β reduces cell viability and strongly impairs BBB functionality.

DISCUSSION

Maintaining BBB integrity is crucial for CNS homeostasis. Hypoxia and ischemia are well known to compromise barrier stability via TJ and extracellular matrix disruption leading to permeability changes (5). Such alterations ultimately contribute to disease progression (4). This study shows that both hypoxia and exogenous TGF β promote cell migration and barrier dysfunction via activation of the proprotein convertase Furin. Blocking Furin abrogates hypoxia and TGF β -induced endothelial migration and permeability indicating a potential future target for vascular-related interventions.

Stimulatory effects of TGF β on cell migration have been reported in intestinal (34) and renal epithelial cells (35), renal microvascular EC (36) and cancer cells (37) under normoxic conditions although high doses of TGF- β may inhibit non-brain EC migration (11, 36). In contrast TGF β did not induce migration of brain ECs in normoxia but only during hypoxic insult in correlation with a study performed on fibroblasts (38). Our observations thus highlight that TGF β signaling activates angiogenic processes in brain EC only during injury conditions, a poignant difference between peripheral cell types and brain EC.

TGF β induces the expression of Furin, a proprotein convertase whose up regulation consistently correlates with tumor cell activation and invasiveness (39). In synovial cells (40) and bovine aortic endothelial cells (41) TGF β was shown to increase Furin transcription. Despite no clear effects on mRNA levels, TGF β treatment clearly induced Furin protein levels in brain EC. Endothelial TGF β can signal through two different TGF β type I receptors, ALK5 and ALK1, which either promote angiogenesis or induce vascular quiescence respectively (11, 32). TGF β signaling through ALK5 activates Smad2 and 3 whilst signaling through ALK1 activates Smad1, 5 and 8 (8, 42). In brain ECs differential Smad signaling was clearly stimuli-related with Smad1/5 phosphorylation being hypoxia-dependent and Smad2 activation being TGF β -dependent. It seems likely that a balance between activation of ALK1 versus ALK5 determines whether ECs remain quiescent or initiate an angiogenic response, an observation previously described by Goumans et al in 2002 (11). In HUVEC and bovine aortic EC ALK5 majorly contributed to angiogenic processes (31, 43). Similarly ALK5 blockade prevented TGF β -induced activation of Smad2 (without affecting pSmad1/5 levels) and suppressed brain EC migration in agreement with those findings. Notably, ALK5 activation was also linked to BBB dysfunction in mouse brain endothelial cell (bEnd.3) monolayers wherein TGF β -induced permeability, via down regulation of the TJ protein Claudin-5, was mediated by Smad3 (44). Overall our data supports the notion that suppressing ALK5 activation

could be key for maintaining BBB function. Our data also suggests Furin modulation is strongly dependent on TGF β /ALK5 signaling and mediated by Smad2 in brain EC in line with other work (18, 40, 45). Whether Smad2 directly or indirectly regulates the effects of TGF β on Furin levels is currently under investigation.

A curious observation from this work is that TGF β induced mature Furin levels under both normoxic and hypoxic conditions but enhanced migration only occurred during hypoxia. Since Furin is a bona fide HIF-1 target (46) it seems plausible that the HIF-1 pathway could somehow contribute to this effect. HIF-1 targeted knockdown studies will further clarify this involvement.

Convertase enzymes exist in an immature preform with a prosegment still attached. Several studies have demonstrated the use of convertase prosegments as inhibitors of their parent enzymes *in vitro* and *ex vivo* (47, 48). During the maturation of Furin, the prosegment acts as a chaperone in the auto-activation pathway as well as a potent auto-inhibitor (49). Indeed profurin overexpression in vascular smooth muscle cells decreased endogenous Furin protein levels and reduced cell proliferation and migration (50). Further hepatic overexpression of the Furin prosegment reduced atherosclerosis progression and decreased vascular remodeling after injury *in vivo* (50). In analogy we observed downregulation of the Furin precursor during hypoxia indicating that potential inhibition of maturation is lifted, resulting in increased Furin activity in line with our migration assay results. This data supports the conclusion that increased Furin activity is ultimately responsible for hypoxic EC migration especially since Furin inhibition not only prevented hypoxic EC migration but also abrogated TGF β -induced effects. In agreement Furin inhibition also attenuated TGF β -enhanced migration of cardiac fibroblasts (51), reduced invasiveness/migration and tumorigenicity of human cancer cells *in vivo* (52) and *in vitro* (47). Hence similar mechanisms underlie brain EC migration and likely contribute to increased barrier permeability during injury conditions. Although not elucidated here, a possible mechanism of Furin-mediated cell movement is likely linked to processing of matrix metalloproteinases (MMPs). Synthesized as inactive propeptides, MMPs undergo endoproteolytic cleavage to attain their active forms (53) resulting in extracellular matrix degradation and barrier disruption (54). Accordingly hypoxia is known to increase Furin-mediated MT1-MMP processing (22, 46). Since Furin inhibition in human cancer cells and cardiac fibroblasts decreased processing and activation of MT1-MMP/MMP-2 (51, 55) it is very tempting to speculate that TGF β -induced brain EC migration is driven by Furin-mediated activation of MMPs.

At the BBB, a TGF β -induced migratory (activated) EC phenotype implies barrier compromise as cytoskeletal alterations lead to a consequential loss of tight junction organization. Indeed TGF β induced brain EC migration and permeability and aggravated injury-induced barrier compromise via an ALK5-dependent mechanism in contrast with a study showing that activation of ALK5 by TGF β inhibits migration in non-brain ECs (13). The unique qualities of BBB ECs including lack of fenestrations, more extensive tight junctions, and reduced pinocytic vesicular transport is clearly significant (6). Although decreased viability of brain EC observed during TGF β treatment was unexpected our data further underscores that negative EC responses to the growth factor can be reversed by inhibiting either ALK5 or Furin activity. Indeed blocking of either abrogated hypoxic barrier disruption agreeing with the notion that these pathways are tightly entwined and induce a migratory/activated phenotype. It is now very evident that Furin activation also plays an important role in injury-induced

BBB disturbance. However an intriguing and unexpected finding, given Furin inhibition prevented short-term hypoxic and TGF β -induced migration, was that combined prolonged TGF β treatment and Furin blockade caused more disruption than TGF β stimulus alone in correlation with dramatically compromised viability. The reason behind this is unclear but disruptive and negative time-dependent effects of sustained TGF β signaling at the BBB and potential confounding effects of modulating multiple downstream pathways must be better understood.

Overall our data suggests both hypoxia- and TGF β -mediated EC permeability can be prevented by blocking Furin and/or ALK5 activation. Notably, a large source of native active TGF β that can be released upon different injury stimuli *in vivo* lies within the perivascular cells and/or the extracellular matrix. Thus further studies using more complex BBB models such as co-culture of ECs with perivascular cells will be needed to provide further insight into the highly complex molecular mechanisms and cellular cross-talk that take place during injury *in vivo*. In summary, controlling Furin or ALK5 activation at the BBB could offer a novel therapeutic option for improving barrier functionality and CNS function during disease.

FIGURE LEGENDS

Fig. 1: TGF β promotes EC migration

Representative micrographs of scratched confluent RBE4 monolayers exposed to 0.4nM TGF β , the ALK5 inhibitor SB431542 (10 μ M) or a combination of both prior to Nx (A) or Hx (B) incubation for 6h. Scale bar=100 μ m. Images were quantified and mean migration in μ m/h is presented (C). Mean \pm SD (n=3), 2way ANOVA, **p<0.01 compared to Nx UNT, #p<0.05 ##p<0.01 ###p<0.001 compared to Hx TGF β .

Fig. 2: Both hypoxia and TGF β modulate Furin

(A) Representative Western blot of Furin precursor (96KDa) and mature (90KDa) forms and HIF-1 α stabilisation at 6h hypoxia with and without TGF β treatment. Blot quantification and graphical representation shows changes of the Furin forms under hypoxia and TGF β treatment (B&C). β -actin was used as loading control. Data are expressed as mean \pm SD of at least n=3 independent experiments. Students t-test and 2way ANOVA, *p<0.05 compared to Nx UNT, §p<0.05 compared to Hx UNT.

Fig. 3: Furin activation is dependent on TGF β receptor-1 ALK5 signaling.

(A) Representative blot showing Furin protein expression after TGF β or SB431542 treatment. Quantitative graphs show comparative levels of both precursor (B) and mature (C) Furin. Mean \pm SD (n=3), Students t-test, §p<0.05, §§p<0.01 compared to Hx UNT, ###p<0.001 to Hx TGF β .

Fig. 4: Furin inhibition prevents EC migration

Representative micrographs of scratched RBE4 monolayers incubated with TGF β , the Furin inhibitors Naphthofluorescein (Naph, 20 μ M) or CMK (20 μ M) or a combination of both prior to Hx exposure for 6h (A). Scale bar =100 μ m. Quantified data (B) shows changes in EC migration under all conditions. Mean \pm SD of at least n=3 independent experiments. 2way ANOVA, ***p<0.001 compared to Nx UNT, ###p<0.01, ##p<0.01 compared to Hx TGF β , §§p<0.01 to Hx UNT.

Fig. 5: TGF β potentiates hypoxia-mediated EC permeability via ALK5

Permeability assay was performed on primary rat endothelial cells grown to confluency on transwells then exposed to 0.4nM TGF β or 10 μ M SB431542 during 24h (A) and 48h (B) hypoxia. Dashed line indicates normoxic baseline measurements. 2way ANOVA and Students t-test, §§p<0.01, §§§p<0.001 compared to Hx CTRL, ##p<0.01 compared to Hx TGF β treatment, Students t-test.

Fig. 6: Furin inhibition prevents hypoxia-induced barrier permeability

Primary rat endothelial cells were grown to confluency on transwells or culture dishes before hypoxic exposure to 0.4nM TGF β , CMK (20 μ M) or CMK combined with TGF β for 48h. Permeability assays (A) and MTT assays (B) were subsequently performed. Mean \pm SD (n=4), 2way ANOVA and Students t-test, **p<0.01, ****p<0.0001 compared to Normoxia, ##p<0.01 to Hx TGF β , §§p<0.01, §§§p<0.001 to Hx UNT.

Supplementary Figure 1: Furin mRNA levels

Quantitative real time PCR shows changes in Furin mRNA levels after treatment with 0.4nM TGF β , 10 μ M ALK5 inhibitor SB431542 or a combination of both prior to 6h

Nx or Hx exposure. Fold changes were calculated with the $\Delta\Delta C_t$ method and normalized to Nx UNT. Mean \pm SD (n=4), 2way ANOVA.

Supplementary Figure 2: TGF β activates Smad2 via ALK5

Representative Western blot (A) and quantification (B) of total and pSmad2 levels after exposure to 0.4nM TGF β , the ALK5 inhibitor (10 μ M) or a combination of both prior to Nx or Hx incubation for 6h. Mean \pm SD (n=4), Students t-test, **p<0.01, ***p<0.001 compared to Nx UNT, #p<0.05 compared to Hx TGF β , °p<0.01 compared to Nx TGF β .

Supplementary Figure 3: Smad1/5 signaling is unaffected by ALK5 blockade

Representative Western blot (A) and quantification (B) of p-Smad1/5 and Smad1 (60kD) protein levels after 6h normoxic or hypoxic exposure with TGF β , ALK5 inhibition or a combination of both (A). Mean \pm SD (n=3), Students t-test, *p<0.05.

Supplementary Figure 4: Furin and ALK5 signaling under normoxia

Representative immunoblot of Furin protein levels after TGF β or SB431542 treatment under normoxic conditions (A). Changes in Furin precursor (B) and mature (C) levels were subsequently quantified. Mean \pm SD (n=4), Students t-test, *p<0.05, **p<0.01 to Nx UNT.

Supplementary Figure 5: Furin inhibition blocks normoxic EC migration

Mean migration (μ m/hr) of normoxic RBE4 cells after exposure to TGF β (0.4nM) or a combination of TGF β and the Furin inhibitors Naphthofluorescein (20 μ M) and CMK (10 μ M). Mean \pm SD (n=3-5), Students t-test, *p<0.05, **p<0.01 to Nx UNT.

ACKNOWLEDGEMENTS

The authors are grateful to Dr. Michele Bernasconi for his scientific input and discussion and Maurizio Roveri for his technical contribution. Part of the experimental work was performed at the Center for Clinical Studies at the Vetsuisse Faculty of the University of Zurich. This work was supported by SNF grant numbers 31003A_150062 and 31003A_170129 to O.O.

REFERENCES

1. Abbott NJ, Patabendige AA, Dolman DE, Yusof SR, Begley DJ. Structure and function of the blood-brain barrier. *Neurobiol Dis.* 2010;37(1):13-25.
2. Brightman MW, Reese TS. Junctions between intimately apposed cell membranes in the vertebrate brain. *J Cell Biol.* 1969;40(3):648-77.
3. Al Ahmad A, Taboada CB, Gassmann M, Ogunshola OO. Astrocytes and pericytes differentially modulate blood-brain barrier characteristics during development and hypoxic insult. *J Cereb Blood Flow Metab.* 2011;31(2):693-705.
4. Zlokovic BV. The blood-brain barrier in health and chronic neurodegenerative disorders. *Neuron.* 2008;57(2):178-201.
5. Engelhardt S, Al-Ahmad AJ, Gassmann M, Ogunshola OO. Hypoxia selectively disrupts brain microvascular endothelial tight junction complexes through a hypoxia-inducible factor-1 (HIF-1) dependent mechanism. *J Cell Physiol.* 2014;229(8):1096-105.
6. Engelhardt S, Huang SF, Patkar S, Gassmann M, Ogunshola OO. Differential responses of blood-brain barrier associated cells to hypoxia and ischemia: a comparative study. *Fluids Barriers CNS.* 2015;12:4.
7. Klempt ND, Sirimanne E, Gunn AJ, Klempt M, Singh K, Williams C, et al. Hypoxia-ischemia induces transforming growth factor beta 1 mRNA in the infant rat brain. *Brain Res Mol Brain Res.* 1992;13(1-2):93-101.
8. Beck K, Schachtrup C. Vascular damage in the central nervous system: a multifaceted role for vascular-derived TGF-beta. *Cell Tissue Res.* 2012;347(1):187-201.
9. Molina F, Rus A, Peinado MA, Del Moral ML. Short-term hypoxia/reoxygenation activates the angiogenic pathway in rat caudate putamen. *J Biosci.* 2013;38(2):363-71.
10. Weiss A, Attisano L. The TGFbeta Superfamily Signaling Pathway. *Wiley Interdisciplinary Reviews: Developmental Biology.* 2013;2(1):47-63.
11. Goumans MJ, Valdimarsdottir G, Itoh S, Rosendahl A, Sideras P, ten Dijke P. Balancing the activation state of the endothelium via two distinct TGF-beta type I receptors. *EMBO J.* 2002;21(7):1743-53.
12. Pardali E, Goumans MJ, ten Dijke P. Signaling by members of the TGF-beta family in vascular morphogenesis and disease. *Trends Cell Biol.* 2010;20(9):556-67.
13. Goumans MJ, Mummery C. Functional analysis of the TGFbeta receptor/Smad pathway through gene ablation in mice. *Int J Dev Biol.* 2000;44(3):253-65.

14. Yang X, Castilla LH, Xu X, Li C, Gotay J, Weinstein M, et al. Angiogenesis defects and mesenchymal apoptosis in mice lacking SMAD5. *Development*. 1999;126(8):1571-80.
15. Blobel GC, Schiemann WP, Lodish HF. Role of transforming growth factor beta in human disease. *N Engl J Med*. 2000;342(18):1350-8.
16. Border WA, Noble NA. Transforming growth factor beta in tissue fibrosis. *N Engl J Med*. 1994;331(19):1286-92.
17. McCaffrey TA, Du B, Consigli S, Szabo P, Bray PJ, Hartner L, et al. Genomic instability in the type II TGF-beta1 receptor gene in atherosclerotic and restenotic vascular cells. *J Clin Invest*. 1997;100(9):2182-8.
18. Blanchette F, Rudd P, Grondin F, Attisano L, Dubois CM. Involvement of Smads in TGFbeta1-induced furin (fur) transcription. *J Cell Physiol*. 2001;188(2):264-73.
19. Seidah NG, Day R, Marcinkiewicz M, Chretien M. Precursor convertases: an evolutionary ancient, cell-specific, combinatorial mechanism yielding diverse bioactive peptides and proteins. *Ann N Y Acad Sci*. 1998;839:9-24.
20. Siegfried G, Basak A, Cromlish JA, Benjannet S, Marcinkiewicz J, Chretien M, et al. The secretory proprotein convertases furin, PC5, and PC7 activate VEGF-C to induce tumorigenesis. *J Clin Invest*. 2003;111(11):1723-32.
21. Siegfried G, Basak A, Prichett-Pejic W, Scamuffa N, Ma L, Benjannet S, et al. Regulation of the stepwise proteolytic cleavage and secretion of PDGF-B by the proprotein convertases. *Oncogene*. 2005;24(46):6925-35.
22. Sato H, Kinoshita T, Takino T, Nakayama K, Seiki M. Activation of a recombinant membrane type 1-matrix metalloproteinase (MT1-MMP) by furin and its interaction with tissue inhibitor of metalloproteinases (TIMP)-2. *FEBS Lett*. 1996;393(1):101-4.
23. Vahatupa M, Cordova ZM, Barker H, Aittomaki S, Uusitalo H, Jarvinen TAH, et al. Furin deficiency in myeloid cells leads to attenuated revascularization in a mouse-model of oxygen-induced retinopathy. *Exp Eye Res*. 2018;166:160-7.
24. Dubois CM, Laprise MH, Blanchette F, Gentry LE, Leduc R. Processing of transforming growth factor beta 1 precursor by human furin convertase. *J Biol Chem*. 1995;270(18):10618-24.
25. Li Y, Liu W, Guan X, Truscott J, Creemers JW, Chen HL, et al. STAT6 and Furin Are Successive Triggers for the Production of TGF-beta by T Cells. *J Immunol*. 2018;201(9):2612-23.
26. Zhang XY, Chang HM, Zhu H, Liu RZ, Leung PCK. BMP6 increases TGF-beta1 production by up-regulating furin expression in human granulosa-lutein cells. *Cell Signal*. 2019;55:109-18.

27. Roux F, Durieu-Trautmann O, Chaverot N, Claire M, Mailly P, Bourre JM, et al. Regulation of gamma-glutamyl transpeptidase and alkaline phosphatase activities in immortalized rat brain microvessel endothelial cells. *J Cell Physiol.* 1994;159(1):101-13.
28. Roll FJ, Madri JA, Albert J, Furthmayr H. Codistribution of collagen types IV and AB2 in basement membranes and mesangium of the kidney. an immunoferritin study of ultrathin frozen sections. *J Cell Biol.* 1980;85(3):597-616.
29. Coisne C, Dehouck L, Faveeuw C, Delplace Y, Miller F, Landry C, et al. Mouse syngenic in vitro blood-brain barrier model: a new tool to examine inflammatory events in cerebral endothelium. *Lab Invest.* 2005;85(6):734-46.
30. Al Ahmad A, Gassmann M, Ogunshola OO. Involvement of oxidative stress in hypoxia-induced blood-brain barrier breakdown. *Microvasc Res.* 2012;84(2):222-5.
31. Jarad M, Kuczynski EA, Morrison J, Vilorio-Petit AM, Coomber BL. Release of endothelial cell associated VEGFR2 during TGF-beta modulated angiogenesis in vitro. *BMC Cell Biol.* 2017;18(1):10.
32. Alsina-Sanchís E, García-Ibáñez Y, Figueiredo AM, Riera-Domingo C, Figueras A, Matias-Guiu X, et al. ALK1 Loss Results in Vascular Hyperplasia in Mice and Humans Through PI3K Activation. *Arteriosclerosis, Thrombosis, and Vascular Biology.* 2018;38(5):1216-29.
33. Coppola JM, Bhojani MS, Ross BD, Rehemtulla A. A small-molecule furin inhibitor inhibits cancer cell motility and invasiveness. *Neoplasia.* 2008;10(4):363-70.
34. Ciacci C, Lind SE, Podolsky DK. Transforming growth factor beta regulation of migration in wounded rat intestinal epithelial monolayers. *Gastroenterology.* 1993;105(1):93-101.
35. Luo D, Guan Q, Wang K, Nguan CYC, Du C. TGF-beta1 stimulates movement of renal proximal tubular epithelial cells in a three-dimensional cell culture via an autocrine TGF-beta2 production. *Exp Cell Res.* 2017;350(1):132-9.
36. Pepper MS. Transforming growth factor-beta: vasculogenesis, angiogenesis, and vessel wall integrity. *Cytokine Growth Factor Rev.* 1997;8(1):21-43.
37. Oft M, Heider KH, Beug H. TGFbeta signaling is necessary for carcinoma cell invasiveness and metastasis. *Curr Biol.* 1998;8(23):1243-52.
38. Mogford JE, Tawil N, Chen A, Gies D, Xia Y, Mustoe TA. Effect of age and hypoxia on TGFbeta1 receptor expression and signal transduction in human dermal fibroblasts: impact on cell migration. *J Cell Physiol.* 2002;190(2):259-65.
39. Khatib AM, Siegfried G, Chretien M, Metrakos P, Seidah NG. Proprotein convertases in tumor progression and malignancy: novel targets in cancer therapy. *Am J Pathol.* 2002;160(6):1921-35.

40. Blanchette F, Day R, Dong W, Laprise MH, Dubois CM. TGFbeta1 regulates gene expression of its own converting enzyme furin. *J Clin Invest.* 1997;99(8):1974-83.
41. Negishi M, Lu D, Zhang YQ, Sawada Y, Sasaki T, Kayo T, et al. Upregulatory expression of furin and transforming growth factor-beta by fluid shear stress in vascular endothelial cells. *Arterioscler Thromb Vasc Biol.* 2001;21(5):785-90.
42. Shi Y, Massague J. Mechanisms of TGF-beta signaling from cell membrane to the nucleus. *Cell.* 2003;113(6):685-700.
43. Li QF, Decker-Rockefeller B, Bajaj A, Pumiglia K. Activation of Ras in the Vascular Endothelium Induces Brain Vascular Malformations and Hemorrhagic Stroke. *Cell Rep.* 2018;24(11):2869-82.
44. McMillin MA, Frampton GA, Seiwel AP, Patel NS, Jacobs AN, DeMorrow S. TGFbeta1 exacerbates blood-brain barrier permeability in a mouse model of hepatic encephalopathy via upregulation of MMP9 and downregulation of claudin-5. *Lab Invest.* 2015;95(8):903-13.
45. Ventura E, Weller M, Burghardt I. Cutting Edge: ERK1 Mediates the Autocrine Positive Feedback Loop of TGF-beta and Furin in Glioma-Initiating Cells. *J Immunol.* 2017;198(12):4569-74.
46. McMahon S, Grondin F, McDonald PP, Richard DE, Dubois CM. Hypoxia-enhanced expression of the proprotein convertase furin is mediated by hypoxia-inducible factor-1: impact on the bioactivation of proproteins. *J Biol Chem.* 2005;280(8):6561-9.
47. Lopez de Cicco R, Bassi DE, Zucker S, Seidah NG, Klein-Szanto AJ. Human carcinoma cell growth and invasiveness is impaired by the propeptide of the ubiquitous proprotein convertase furin. *Cancer Res.* 2005;65(10):4162-71.
48. Zhong M, Munzer JS, Basak A, Benjannet S, Mowla SJ, Decroly E, et al. The prosegments of furin and PC7 as potent inhibitors of proprotein convertases. In vitro and ex vivo assessment of their efficacy and selectivity. *J Biol Chem.* 1999;274(48):33913-20.
49. Thomas G. Furin at the cutting edge: from protein traffic to embryogenesis and disease. *Nat Rev Mol Cell Biol.* 2002;3(10):753-66.
50. Lei X, Basu D, Li Z, Zhang M, Rudic RD, Jiang XC, et al. Hepatic overexpression of the prodomain of furin lessens progression of atherosclerosis and reduces vascular remodeling in response to injury. *Atherosclerosis.* 2014;236(1):121-30.
51. Stawowy P, Margeta C, Kallisch H, Seidah NG, Chretien M, Fleck E, et al. Regulation of matrix metalloproteinase MT1-MMP/MMP-2 in cardiac fibroblasts by TGF-beta1 involves furin-convertase. *Cardiovasc Res.* 2004;63(1):87-97.

52. Jaaks P, D'Alessandro V, Grob N, Buel S, Hajdin K, Schafer BW, et al. The Proprotein Convertase Furin Contributes to Rhabdomyosarcoma Malignancy by Promoting Vascularization, Migration and Invasion. *PLoS One*. 2016;11(8):e0161396.
53. Murphy G, Stanton H, Cowell S, Butler G, Knauper V, Atkinson S, et al. Mechanisms for pro matrix metalloproteinase activation. *APMIS*. 1999;107(1):38-44.
54. Candelario-Jalil E, Yang Y, Rosenberg GA. Diverse roles of matrix metalloproteinases and tissue inhibitors of metalloproteinases in neuroinflammation and cerebral ischemia. *Neuroscience*. 2009;158(3):983-94.
55. Bassi DE, Lopez De Cicco R, Mahloogi H, Zucker S, Thomas G, Klein-Szanto AJ. Furin inhibition results in absent or decreased invasiveness and tumorigenicity of human cancer cells. *Proc Natl Acad Sci U S A*. 2001;98(18):10326-31.


Figure 1: TGF β promotes EC migration


Figure 2: Both hypoxia and TGF β modulate Furin

Figure 3: Furin activation is dependent on ALK5 signaling




Figure 4: Furin inhibition prevents EC migration






Figure 5: TGF β potentiates hypoxia-mediated EC permeability via ALK5

Figure 6: Furin inhibition prevents hypoxia-induced barrier permeability



AUTHOR CONFLICT OF INTEREST STATEMENT (for ECR-19-511)

We wish to confirm that there are no known conflicts of interest associated with this publication and there has been no significant financial support for this work that could have influenced its outcome.

We confirm that the manuscript has been read and approved by all named authors and that there are no other persons who satisfy the criteria for authorship but are not listed. We further confirm that the order of authors listed in the manuscript has been approved by all of us.

We confirm that we have given due consideration to the protection of intellectual property associated with this work and that there are no impediments to publication, including the timing of publication, with respect to intellectual property. In so doing we confirm that we have followed the regulations of our institutions concerning intellectual property.

We further confirm that any aspect of the work covered in this manuscript that has involved either experimental animals or human patients has been conducted with the ethical approval of all relevant bodies and that such approvals are acknowledged within the manuscript.

We understand that the Corresponding Author is the sole contact for the Editorial process (including Editorial Manager and direct communications with the office). He/she is responsible for communicating with the other authors about progress, submissions of revisions and final approval of proofs. We confirm that we have provided a current, correct email address that is accessible by the Corresponding Author.

Signed and dated by all authors below:

Julia Baumann	 24.6.19
Sheng-Fu Huang	 24.06.19
Max Gassmann	 25.6.19
Chih-Chieh Tsao	 24.06.19
Omolara Ogunshola	 24-6-19